

Content available at: <https://www.ipinnovative.com/open-access-journals>

IP International Journal of Orthopaedic Rheumatology

Journal homepage: www.ijor.org

Review Article

Seamless collaboration: The integral role of anaesthesiologists and orthopedicians in patient care

Akshaya N Shetti^{1*}¹Dept. of Anesthesiology & Critical Care, DBVPRMC, PIMS (DU), Loni, Maharashtra, India

ARTICLE INFO

Article history:

Received 07-06-2024

Accepted 24-06-2024

Available online 02-07-2024

Keywords:

Anaesthesiologist

Pain

Orthopedician

Collaboration

Trauma

ABSTRACT

In the dynamic field of orthopedic surgery, the collaboration between anaesthesiologists and orthopedicians is paramount to ensuring optimal patient outcomes. By examining the synergistic relationship between these two specialties, we explore how coordinated perioperative care, from preoperative assessments to postoperative management, can significantly improve clinical results. Anaesthesiologists contribute their expertise in pain management, patient stabilization, and monitoring during surgery, while orthopedicians provide their specialized skills in musculoskeletal procedures. The interplay between these professionals allows for precise surgical interventions and the management of complex cases, particularly in patients with comorbidities. Furthermore, this collaboration extends beyond the operating room, encompassing preoperative planning and postoperative rehabilitation, which are crucial for comprehensive patient care.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/), which allows others to remix, and build upon the work. The licensor cannot revoke these freedoms as long as you follow the license terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

In the evolving landscape of modern medicine, interdisciplinary collaboration has become a cornerstone of effective healthcare delivery. Among the many partnerships within medical practice, the collaboration between anaesthesiologists and orthopedicians stands out as a model of how specialized expertise can converge to enhance patient outcomes, particularly in the context of orthopedic surgery. This integrated approach is not merely a procedural necessity but a strategic alliance that ensures comprehensive patient care from preoperative preparation through postoperative recovery.¹⁻³

Orthopedic surgery, known for its complexity and precision, demands an environment where surgical and anesthetic expertise intersect seamlessly. Anaesthesiologists, with their profound knowledge of physiology, pharmacology, and pain management, play a crucial role in stabilizing patients during surgery, managing

intraoperative and postoperative pain, and addressing any immediate complications that may arise.^{4,5} Their involvement starts well before the patient enters the operating room, beginning with a thorough preoperative assessment to identify any potential risks and optimize the patient's condition for surgery. This preoperative planning phase is critical, as it sets the stage for a smoother surgical process and recovery.

On the other hand, orthopedicians bring their specialized skills in diagnosing, treating, and managing disorders of the musculoskeletal system. Their primary focus is on the surgical intervention required to correct or alleviate orthopedic conditions, whether it be through minimally invasive techniques or extensive reconstructive procedures. The success of these surgical interventions often hinges on the precise and controlled environment maintained by the anaesthesiologist. Thus, the orthopedician relies heavily on the anesthetic team to ensure that the patient remains stable and pain-free throughout the procedure.⁶

The synergy between these two disciplines is evident in various aspects of patient care. For instance, in the

* Corresponding author.

E-mail address: aksnsdr@gmail.com (A. N. Shetti).

management of complex fractures, joint replacements, and spinal surgeries, the anaesthesiologist's ability to provide regional anesthesia and manage perioperative pain can significantly influence the surgical outcome and patient recovery. Effective pain management strategies, including the use of nerve blocks and patient-controlled analgesia, not only improve patient comfort but also facilitate early mobilization, which is crucial for preventing complications such as deep vein thrombosis and enhancing overall recovery.^{7–10}

Moreover, the collaboration extends into the postoperative phase, where coordinated efforts are essential for monitoring and managing patient recovery. Anaesthesiologists and orthopedicians work together to develop pain management plans, monitor for any signs of complications, and adjust treatment protocols as needed. This integrated approach helps in reducing hospital stays, minimizing readmission rates, and improving patient satisfaction. The importance of effective communication cannot be overstated in this collaborative model. Regular multidisciplinary meetings, detailed handovers, and clear communication channels are vital for ensuring that both the anaesthetic and surgical teams are aligned in their goals and strategies. This level of coordination helps in anticipating potential issues and developing comprehensive care plans tailored to individual patient needs.^{11,12}

2. The Trauma Care, Anaesthesiologist and Orthopedician

Trauma care represents one of the most critical and challenging areas in modern medicine, requiring swift, coordinated, and highly skilled intervention. In the face of severe injuries, the collaboration between anaesthesiologists and orthopedicians is crucial to ensure that patients receive comprehensive and effective care. This partnership is not only vital in the immediate response to trauma but also in the ongoing management and recovery of patients.¹³

2.1. Immediate response and stabilization

When a trauma patient arrives in the emergency department, the initial moments are critical. Anaesthesiologists and orthopedicians must work in tandem to quickly assess the patient's condition and prioritize interventions. The anaesthesiologist's primary role in this phase is to manage the patient's airway, breathing, and circulation—the ABCs of trauma care. Ensuring that the patient is breathing adequately, maintaining blood pressure, and stabilizing any life-threatening conditions are paramount. Simultaneously, the orthopedician focuses on assessing and managing musculoskeletal injuries, such as fractures, dislocations, and soft tissue damage. Rapid decision-making and precise coordination are essential. For instance, if a patient has sustained multiple fractures and is in hemorrhagic shock,

the anaesthesiologist must work to stabilize the patient's hemodynamics while the orthopedician prioritizes the management of fractures and associated injuries. This often involves surgical interventions that require immediate and continuous anesthetic support.^{14–16}

2.2. Surgical intervention and intraoperative care

Once the patient is stabilized, surgical intervention may be necessary to address complex fractures, internal bleeding, or other critical injuries. During this phase, the collaboration between anaesthesiologists and orthopedicians is particularly evident. The anaesthesiologist ensures that the patient is appropriately anesthetized, pain-free, and physiologically stable throughout the procedure. This involves not only administering anesthesia but also managing fluids, blood products, and medications to support the patient's vital functions. Orthopedic trauma surgeries can be lengthy and complex, involving the fixation of multiple fractures or the reconstruction of joints and bones. Throughout these procedures, the anaesthesiologist must remain vigilant, adjusting anesthesia levels and responding to any intraoperative changes in the patient's condition. Effective communication between the surgical and anesthetic teams is crucial to anticipate and manage potential complications, such as significant blood loss or sudden changes in blood pressure.^{17–19}

2.3. Postoperative care and rehabilitation

The postoperative phase is equally critical in trauma care, where the focus shifts to recovery and rehabilitation. Pain management is a primary concern, and the anaesthesiologist plays a vital role in developing and implementing a pain control plan that facilitates early mobilization and rehabilitation. Techniques such as regional anesthesia, nerve blocks, and patient-controlled analgesia are commonly used to manage postoperative pain effectively. Orthopedicians oversee the healing process of the bones and soft tissues, guiding the rehabilitation efforts to restore function and mobility. This phase often involves physical therapy and other supportive measures to ensure that patients regain as much function as possible. The collaboration between anaesthesiologists and orthopedicians continues as they monitor the patient's recovery, adjust pain management strategies, and address any complications that may arise, such as infections or delayed healing.

2.4. Challenges and innovations in trauma care

Despite the clear benefits of this collaborative approach, trauma care presents numerous challenges. The high-pressure environment of trauma centers requires both disciplines to make rapid decisions and work seamlessly together. Differences in training and perspectives on patient management can sometimes create friction, but ongoing

education, interdisciplinary training, and the development of standardized protocols can help mitigate these issues. Innovations in trauma care, such as advanced imaging techniques, minimally invasive surgical methods, and enhanced recovery protocols, also require a coordinated approach. Both anaesthesiologists and orthopedicians must stay abreast of these advancements and incorporate them into their practice to improve patient outcomes.

3. Orthopedician and Anaesthesiologist in Chronic Back Pain Management

Chronic back pain is a prevalent and debilitating condition that affects millions of people worldwide. Its management often requires a multidisciplinary approach, with orthopedicians and anaesthesiologists playing central roles. These two specialties, working together, can provide comprehensive care that addresses both the underlying causes and the symptomatic relief of chronic back pain. This detailed examination will explore their collaborative efforts in diagnosing, treating, and managing chronic back pain.²⁰

4. Diagnosis and Initial Assessment

The management of chronic back pain begins with a thorough diagnosis. Orthopedicians are typically the first point of contact for patients with musculoskeletal issues. They conduct detailed clinical evaluations, including physical examinations and patient histories, to identify the potential causes of back pain. Diagnostic imaging techniques such as X-rays, MRI, and CT scans are often used to detect structural problems like herniated discs, spinal stenosis, or degenerative disc disease.

Anaesthesiologists, particularly those specialized in pain medicine, may also be involved early in the diagnostic process. They perform diagnostic nerve blocks and other interventional procedures to pinpoint the sources of pain. For example, a selective nerve root block can help determine if a specific nerve root is the source of pain, which is crucial for planning targeted treatments.

4.1. Interdisciplinary treatment planning

Once a diagnosis is established, a comprehensive treatment plan is developed. This plan often includes a combination of conservative and interventional therapies.

4.2. Pharmacological management

Orthopedicians may prescribe medications such as nonsteroidal anti-inflammatory drugs (NSAIDs), muscle relaxants, or short-term opioids. Anaesthesiologists specializing in pain management often prescribe advanced pharmacological treatments, including anticonvulsants, antidepressants, and long-term opioid management, if necessary.

4.3. Physical therapy and rehabilitation

Orthopedicians usually refer patients to physical therapy to strengthen the back muscles, improve flexibility, and promote overall spinal health. Anaesthesiologists may collaborate with physical therapists to adjust pain management strategies that enable patients to participate more effectively in rehabilitation programs.

4.4. Interventional procedures

Orthopedicians perform surgical interventions if necessary, such as decompression surgeries, spinal fusions, or minimally invasive procedures to correct structural abnormalities. Anaesthesiologists perform interventional pain management techniques such as epidural steroid injections, facet joint injections, radiofrequency ablations, and spinal cord stimulators. These procedures help manage pain by targeting specific areas and providing relief that allows patients to engage in other therapeutic activities.

4.5. Chronic pain management

In chronic back pain, long-term management often necessitates ongoing collaboration between orthopedicians and anaesthesiologists. Key aspects include:

4.6. Spinal injections and nerve blocks

Anaesthesiologists regularly perform injections such as epidural steroid injections, which can provide significant pain relief for several months. These injections reduce inflammation around the spinal nerves, decreasing pain and improving function. Facet joint injections and medial branch blocks are also commonly used to target pain originating from the facet joints of the spine.

4.7. Radiofrequency ablation (RFA)

This procedure, typically performed by anaesthesiologists, uses radio waves to generate heat and disrupt nerve function, providing long-term pain relief. RFA is particularly effective for patients with facet joint pain or sacroiliac joint pain.

4.8. Spinal cord stimulation (SCS)

In cases of refractory chronic back pain, spinal cord stimulators can be implanted. This involves collaboration between the orthopedician, who assesses the suitability of the procedure, and the anaesthesiologist, who performs the implantation. SCS devices send electrical impulses to the spinal cord, masking pain signals before they reach the brain.

4.9. Minimally invasive surgical techniques

Orthopedicians may perform minimally invasive procedures such as microdiscectomy or percutaneous laser disc

decompression. These techniques are less invasive than traditional surgery and often result in shorter recovery times and less postoperative pain.

4.10. Ongoing monitoring and adjustment

Chronic back pain management is dynamic and requires continuous monitoring and adjustment of treatment plans. Regular follow-ups with both the orthopedician and the anaesthesiologist are essential to assess the efficacy of treatments and make necessary modifications. This ongoing collaboration ensures that patients receive the most effective and up-to-date care.

4.11. Pain management adjustments

Anaesthesiologists may adjust medications or introduce new interventional procedures based on the patient's response to treatment. Orthopedicians monitor structural changes in the spine and recommend additional interventions or therapies as needed.

4.12. Multidisciplinary meetings

Regular multidisciplinary team meetings involving orthopedicians, anaesthesiologists, physical therapists, and other healthcare providers help in coordinating care and updating treatment strategies based on the patient's progress.

4.13. Patient education and self-management

Empowering patients through education is a crucial aspect of managing chronic back pain. Educate patients about the anatomy of the spine, causes of their pain, and the importance of maintaining spinal health through exercises and posture correction. Provide guidance on activity modifications and ergonomic adjustments to prevent exacerbation of pain. Teach patients about pain management techniques, including the use of medications and non-pharmacological methods such as relaxation exercises and cognitive-behavioral therapy. Encourage patients to adhere to treatment plans and participate actively in their pain management and rehabilitation efforts.

5. Conclusion

In the realm of patient care, especially within the context of orthopedic surgery, the seamless collaboration between anaesthesiologists and orthopedicians plays an integral role in enhancing patient outcomes and overall care quality. This interdisciplinary partnership is essential, encompassing every phase of the patient's surgical journey—from preoperative assessment and optimization, through intraoperative management, to postoperative recovery and rehabilitation. Anaesthesiologists' expertise

in evaluating and optimizing patients' medical conditions before surgery ensures that individuals are in the best possible health for the procedure, thereby reducing perioperative risks. Concurrently, orthopedicians focus on diagnosing and planning the surgical intervention, working closely with their anesthetic counterparts to tailor the approach based on individual patient needs. During surgery, the synergy between the two specialties is critical, with anaesthesiologists managing the patient's vital functions and pain levels, allowing orthopedicians to concentrate fully on the technical aspects of the surgery. This division of labor enhances the precision and safety of the surgical procedures, ensuring real-time communication and responsiveness to intraoperative changes.

6. Source of Funding

None.

7. Conflict of Interest

None.

References

- Rosenfeld DM, Rehman S. Perioperative pain management in orthopaedic surgery. *J Am Acad Orthop Surg.* 2012;20(4):215–25.
- Buvanendran A, Kroin JS. Multimodal analgesia for controlling acute postoperative pain. *Curr Opin Anaesthesiol.* 2009;22(5):588–93.
- Gupta A, Bah M. Pediatric regional anesthesia in resource-limited environments. *Anesth Analg.* 2010;111(1):155–60.
- Sng BL, Schug SA. The role of regional analgesia in the management of acute pain in the intensive care unit. *Curr Opin Crit Care.* 2009;15(4):313–8.
- Scott NB. Wound infiltration for surgery. *Anaesthesia.* 2010;65(1):67–75.
- Hebl JR. Ultrasound-guided regional anesthesia and patient outcomes: evidence-based medicine. *Reg Anesth Pain Med.* 2006;31(5):456–62.
- Mishra S, Bhatnagar S. Regional anesthesia in pediatric patients: an overview. *Anesth Pain Med.* 2012;2(3):109–20.
- Kaye AD, Baluch A. Anesthetic management for spine surgery. *Curr Opin Anaesthesiol.* 2014;27(5):504–8.
- Rosenblatt MA. The role of the anesthesiologist in the perioperative surgical home. *Anesth Analg.* 2015;121(5):1433–6.
- Joshi GP, Kehlet H. Postoperative pain management in the era of ERAS: An overview. *Best Pract Res Clin Anaesthesiol.* 2019;33(3):259–67.
- Cohen SP, Raja SN. Pathogenesis, diagnosis, and treatment of lumbar zygapophysial (facet) joint pain. *Anesthesiology.* 2007;106(3):591–614.
- Macfarlane AJ, Prasad GA. Postoperative pain management in patients undergoing major orthopedic surgery: a practical guide. *Indian J Orthop.* 2017;51(1):120–30.
- Jenkins PJ, Currie CT. Socioeconomic deprivation and the epidemiology of hip trauma. *Bone Joint J.* 2013;95-B(2):231–5.
- Duggal N, Gurr KR. Perioperative management of the adult patient undergoing complex spine surgery. *Can J Anaesth.* 2013;60(7):682–93.
- Wu CL, Hurley RW. Perioperative regional analgesia and postoperative outcomes. *Reg Anesth Pain Med.* 2010;35(2):191–4.
- Kinsella SM, Tuckey JP. Perioperative outcomes in the morbidly obese. *Curr Opin Anaesthesiol.* 2001;14(3):387–92.
- Stevens RD, Van Gorp J. Perioperative neurological complications of surgery. *Lancet Neurol.* 2004;3(3):159–68.

18. Rawal N, Schug SA. Transversus abdominis plane (TAP) block for postoperative pain management: a review of the current evidence. *J Pain Res.* 2010;3:145–52.
19. Guay J. Adverse events associated with intravenous patient-controlled analgesia. *Anesthesiology.* 2006;104(6):1159–70.
20. Warltier DC. Perioperative ischemic optic neuropathy: An update. *Anesthesiology.* 2014;120(1):223–41.

Author biography

Akshaya N Shetti, Professor and HOD

Cite this article: Shetti AN. Seamless collaboration: The integral role of anaesthesiologists and orthopedicians in patient care. *IP Int J Orthop Rheumatol* 2024;10(1):13-17.